



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the “Company”), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

Note: Stereotactic Radiosurgery and Stereotactic Body Radiotherapy is addressed separately in medical policy 00045.

Note: The use of Radiofrequency Ablation (RFA) of Primary or Metastatic Liver Tumors is addressed separately in medical policy 00182.

Note: Cryosurgical Ablation of Primary or Metastatic Liver Tumors is addressed separately in medical policy 00220.

When Services Are Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member’s contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Osteoid Osteomas

Based on review of available data, the Company may consider radiofrequency ablation (RFA) of osteoid osteomas to be **eligible for coverage.****

Osteolytic Bone Metastases

Based on review of available data, the Company may consider radiofrequency ablation (RFA) of painful bony metastases in individuals who have failed or who are poor candidates for standard treatments such as opioids or radiation therapy to be **eligible for coverage.****

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

Localized Renal Malignancy

Based on review of available data, the Company may consider radiofrequency ablation (RFA) for clinically localized, suspected renal malignancy for individuals with peripheral lesions that are less than or equal to 4 cm in diameter to be **eligible for coverage**.**

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Non-Small Cell Lung Cancer

Based on review of available data, the Company may consider radiofrequency ablation (RFA) of non-small cell lung cancer (NSCLC) to be **eligible for coverage**.**

Patient Selection Criteria

Coverage eligibility for RFA of NSCLC will be considered when **ALL** of the following criteria are met:

- Surgical or radiation treatment with curative intent is considered appropriate based on stage of disease, however medical co-morbidity renders the individual unfit for those interventions; **AND**
- No tumor has a maximum diameter of greater than 3.0 cm; **AND**
- Tumors are located at least 1 cm from the trachea, main bronchi, esophagus, aorta, aortic arch branches, pulmonary artery and the heart.

Metastatic Malignant Tumor(s) to the Lung

Based on review of available data, the Company may consider radiofrequency ablation (RFA) to treat malignant nonpulmonary tumor(s) metastatic to the lung to be **eligible for coverage**.**

Patient Selection Criteria

Coverage eligibility for RFA to treat malignant nonpulmonary tumor(s) metastatic to the lung will be considered when **ALL** of the following criteria are met:

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

- When it is necessary to preserve lung function because surgical resection or radiotherapy is likely to worsen pulmonary status substantially, OR when the individual is not considered a surgical candidate; **AND**
- There is no evidence of extra-pulmonary metastases; **AND**
- No more than 3 tumors per lung should be ablated; **AND**
- No tumor has a maximum diameter greater than 3.0 cm; **AND**
- Tumors are located at least 1 cm from the trachea, main bronchi, esophagus, aorta, aortic arch branches, pulmonary artery and the heart; **AND**
- If a repeat procedure, at least 12 months have elapsed since the prior ablation.

When Services Are Considered Investigational

Coverage is not available for investigational medical treatments or procedures, drugs, devices or biological products.

Based on review of available data, the Company considers radiofrequency ablation (RFA) of tumors outside the liver to be **investigational*** when the above criteria are not met and for all other tumors, including but not limited to:

- Breast, the head and neck, thyroid, pancreas, adrenal gland, ovary, and pelvic/abdominal metastases of unspecified origin.

Background/Overview

Health Disparities in Certain Solid Tumor Types

Based on data from 2014 through 2018, age-adjusted breast cancer mortality is approximately 40% higher among Black women compared to non-Hispanic White women in the United States (27.7 vs 20.0 deaths per 100,000 women), despite a lower overall incidence of breast cancer among Black women (125.8 vs 139.2 cases per 100,000 women). Experts postulate that this divergence in mortality may be related to access issues— Black women are more likely than White women to lack health insurance, limiting access to screening and appropriate therapies. Socioeconomic status is also a driver in health and health outcome disparities related to breast cancer. Women with low incomes have significantly lower rates of breast cancer screening, a higher probability of late-stage diagnosis, and are less likely to receive high-quality care, resulting in higher mortality from breast cancer.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

Based on data from 2016 through 2020, kidney cancer is more common in men than women and occurs more often in American Indian and Alaskan Native individuals, followed by Black and Hispanic individuals. American Indians and Alaska Natives have higher death rates from kidney cancer than any other racial or ethnic group. A cohort study by Howard et al (2021) included 158,445 patients with localized kidney cancer from the National Cancer Database between 2010 and 2017. Investigators found that female patients were treated more aggressively compared with male patients, with lower adjusted odds of undertreatment and higher adjusted odds of overtreatment. They also found that Black and Hispanic patients had higher adjusted odds of undertreatment and overtreatment compared to White patients, and uninsured status was associated with lower adjusted odds of overtreatment and higher adjusted odds of undertreatment. These results suggest that sex, race and ethnicity, and socioeconomic status are associated with disparities in guideline-based treatment for localized kidney cancer, specifically, with increased rates of non-guideline-based treatment for women and Black and Hispanic patients.

Radiofrequency Ablation

Radiofrequency ablation (RFA) was initially developed to treat inoperable tumors of the liver (see medical policy 00182). Recently, studies have reported on the use of RFA to treat other tumors. For some of these, RFA is being investigated as an alternative to surgery for operable tumors. Well-established local or systemic treatment alternatives are available for each of these malignancies. The hypothesized advantages of RFA for these cancers include improved local control and those common to any minimally invasive procedure (eg, preserving normal organ tissue, decreasing morbidity, decreasing length of hospitalization).

Goals of RFA may include (1) controlling local tumor growth and preventing recurrence; (2) palliating symptoms; and (3) extending survival duration for patients with certain tumors. The effective volume of RFA depends on the frequency and duration of applied current, local tissue characteristics, and probe configuration (eg, single vs. multiple tips). RFA can be performed as an open surgical procedure, laparoscopically or percutaneously, with ultrasound or computed tomography guidance.

Potential complications associated with RFA include those caused by heat damage to normal tissue adjacent to the tumor (eg, intestinal damage during RFA of kidney), structural damage along the probe track (eg, pneumothorax as a consequence of procedures on the lung), and secondary tumors (if cells seed during probe removal).

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

The U.S. Food and Drug Administration (FDA) issued a statement in September 2008, concerning the regulatory status of RFA. The FDA has cleared RFA devices for the general indication of soft tissue cutting, coagulation, and ablation by thermal coagulation necrosis. Under this general indication, RFA can be used to ablate tumors, including lung tumors. Some RFA devices have been cleared for additional specific treatment indications, including partial or complete ablation of nonresectable liver lesions and palliation of pain associated with metastatic lesions involving bone. The FDA has not cleared any RFA devices for the specific treatment indication of partial or complete ablation of lung tumors, citing lack of sufficient clinical data to establish safety and effectiveness for this purpose. The FDA has received reports of death and serious injuries associated with the use of RFA devices in the treatment of lung tumors.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to federal regulations, other plan medical policies, and accredited national guidelines.

Description

In radiofrequency ablation (RFA), a probe is inserted into the center of a tumor; then, prong-shaped, non-insulated electrodes are projected into the tumor. Next, heat is generated locally by an alternating, high-frequency current that travels through the electrodes. The localized heat treats the tissue adjacent to the probe, resulting in a 3 cm to 5.5 cm sphere of dead tissue. The cells killed by RFA are not removed but are gradually replaced by fibrosis and scar tissue. If there is a local recurrence, it occurs at the edge and can sometimes be retreated. RFA may be performed percutaneously, laparoscopically, or as an open procedure.

Summary of Evidence

For individuals who have painful osteolytic bone metastases who have failed or are poor candidates for standard treatments who receive RFA, the evidence includes a prospective cohort study and case series. Relevant outcomes are symptoms, change in disease status, quality of life (QOL), medication

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

use, and treatment-related morbidity. A prospective cohort study and case series have shown clinically significant pain relief (defined as a decrease of 2 units from baseline on the Brief Pain Inventory scale) or reduction in opioid use following treatment of painful osteolytic metastases. A multicenter, prospective study reported significant reductions in pain through the 6-month follow-up period, with 59% of patients achieving immediate improvement in pain within 3 days of RFA. The population is comprised of patients with few or no treatment options, for whom short-term pain relief is an appropriate clinical outcome. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have painful osteoid osteomas who receive RFA, the evidence includes numerous observational studies and systematic reviews of these studies. Relevant outcomes are symptoms, change in disease status, QOL, medication use, and treatment-related morbidity. In a systematic review of thermal ablation techniques, clinical success (pain-free) was achieved in 94% to 98% of patients. Most patients (89% to 96%) remained pain-free when assessed during longer-term follow-up. Another systematic review reported similar success rates noting an average 8.3% failure rate among patients receiving computed tomography-guided RFA. Although no randomized trials of RFA for osteoid osteomas have been performed, the uncontrolled studies have demonstrated RFA can provide adequate symptom relief with minimal complications, for a population for whom short-term symptom relief and avoidance of invasive procedures are appropriate clinical outcomes. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have localized renal cell carcinoma (RCC) that is no more than 4 cm in size who receive RFA, the evidence includes a randomized controlled trial (RCT), numerous observational studies, and systematic reviews of these studies. Relevant outcomes are overall survival (OS), change in disease status, QOL, and treatment-related morbidity. A recent meta-analysis that included only an RCT and cohort studies found that RFA was as effective as nephrectomy for small renal tumors, with a reduction in complications. Another recent meta-analysis found that partial nephrectomy (PN) was superior to ablative techniques (the study included RFA but also cryoablation and microwave ablation) in overall mortality and local recurrence but not in cancer-specific mortality. It also found fewer complications and improved renal function with ablation. A meta-analysis from 2022 found that PN was superior to ablation (RFA, cryoablation, and microwave ablation) in local recurrence. Overall complications, decline in renal function, and cancer-specific mortality rates did not differ between ablation and PN. Although inconsistent, the

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

evidence does suggest that, for small renal tumors, RFA may result in a similar rate of disease progression with a lower complication rate than nephrectomy. However, comparative trials are needed to determine with greater certainty the effects of these treatments in the same patient population. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have inoperable primary pulmonary tumors or nonpulmonary tumors metastatic to the lung who receive RFA, the evidence includes prospective observational studies and systematic reviews of these studies. Relevant outcomes are OS, change in disease status, QOL, and treatment-related morbidity. A multicenter study found that for tumors less than 3.5 cm in size, RFA can lead to a complete response in as many as 88% of patients for at least 1 year. Two-year survival rates have been reported to range from 41% to 75% in case series, with 5-year survival rates of 20% to 27%. In general, the evidence suggests that RFA results in adequate survival and tumor control in patients who are not surgical candidates, with low morbidity rates. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have breast tumors who receive RFA, the evidence includes observational studies and systematic reviews of these studies. Relevant outcomes are OS, change in disease status, QOL, and treatment-related morbidity. Evidence has reported varied and incomplete ablation rates with concerns about postablation tumor cell viability. Long-term improvements in health outcomes have not been demonstrated. Additionally, available studies do not permit comparisons with conventional breast-conserving procedures. Further prospective studies, with long-term follow-up, should focus on whether RFA of the breast for small tumors can provide local control and survival rates compared with conventional breast-conserving treatment. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have benign thyroid tumors who receive RFA, the evidence includes RCTs, prospective studies, case series, and systematic reviews of these studies. Relevant outcomes are symptoms, change in disease status, QOL, medication use, and treatment-related morbidity. Systematic reviews have demonstrated that RFA results in a significant reduction in thyroid nodule size, with a 2020 review showing that these changes remain durable through at least 36 months. Complication rates are generally low, but include voice changes. The data are limited by significant heterogeneity in meta-analyses, a lack of generalizability to populations outside Republic of Korea and Italy, and a lack of comparators more relevant to practice in the United States. Further studies

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

comparing RFA to percutaneous ethanol injection (PEI) or surgery would be more informative in determining the potential utility of RFA in patients with symptomatic or large benign thyroid tumors as these are the recommended treatment options per the American Thyroid Association. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have miscellaneous tumors (eg, head and neck, thyroid cancer, pancreas) who receive RFA, the evidence includes a few case series, prospective observational studies, and retrospective comparative studies. Relevant outcomes are OS, change in disease status, QOL, and treatment-related morbidity. There is a limited evidence base for these tumor types. Reporting on outcomes or comparisons with other treatments is limited. These studies do not permit conclusions on the health benefits of RFA. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

Additional Information

Clinical input obtained in 2010 supported use of RFA for localized RCC that is no more than 4 cm in size when preservation of kidney function is necessary, and a standard surgical approach is likely to worsen kidney function substantially or when the patient is not considered a surgical candidate. Thus, absent other treatment options, RFA for small renal cell tumors was judged to be medically necessary.

Supplemental Information

Clinical Input From Physician Specialty Societies and Academic Medical Centers

2010 Input

In response to requests, input was received from 2 physician specialty societies (4 reviewers) and 2 academic medical centers (4 reviewers) while this policy was under review in 2010. Input was similar to that received in 2009, except support for the use of radiofrequency ablation (RFA) to treat lung tumors was declined (only 1 respondent indicated this was an option in tumors metastatic to lung). One respondent also indicated a potential use for adrenal tumors. Input supported RFA for localized renal cell carcinoma no more than 4 cm in size when preservation of kidney function is necessary and a standard surgical approach would likely substantially worsen kidney function or when the patient is not considered a surgical candidate.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

2009 Input

In response to requests, input was received from 1 physician specialty society (4 reviews) and from 2 academic medical centers (3 reviews) while this policy was under review in 2009. All reviewers supported the use of RFA in the treatment of painful bone metastases that have failed standard treatment and in the treatment of osteoid osteomas. Reviewers were divided over the use of RFA for lung tumors, although several agreed that, while it may be useful in a select population of patients, it should be used in the clinical trial setting. Reviewers were also split with regard to RFA in the treatment of renal tumors, with some supporting its use in a select population of patients. With the exception of 1 disagreement and 1 nonresponse, the reviewers agreed to the investigational statement on the use of RFA in all other tumors outside the liver that are addressed in this policy.

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in 'Supplemental Information' if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Chest Physicians

The American College of Chest Physicians (2013) guidelines on the treatment of stage I and II non-small-cell lung cancer (NSCLC) have indicated RFA has been used effectively in clinical stage I NSCLC. Therefore, in medically inoperable patients, peripheral NSCLC tumors less than 3 cm may be treated with RFA. The College also collaborated with the Society of Thoracic Surgeons to develop consensus guidelines on the treatment of high-risk patients with stage I NSCLC. These 2012 consensus guidelines indicated RFA is an alternative treatment option for patients who are not surgical candidates due to severe medical comorbidity.

American Head and Neck Society - Endocrine Surgery Section

An international, multidisciplinary consensus statement on RFA and related ultrasound-guided ablation technologies for the treatment of benign and malignant thyroid disease was released in 2022 through a collaboration of international professional societies, including the Endocrine Surgery Section of the American Head and Neck Society. Select relevant recommendations from the guideline are listed in Table 1.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

Table 1. Summary of RFA Recommendations for Treatment of Benign and Malignant Thyroid Disease*

Recommendation 1	US-guided ablation procedures may be used as a first-line alternative to surgery for patients with benign thyroid nodules contributing to compressive and/or cosmetic symptoms.
Recommendation 2	Although less efficacious than surgery or RAI in normalizing thyroid function, thermal ablation procedures can be a safe therapeutic alternative in patients with an autonomously functional thyroid nodule and contraindications to first-line techniques.
Recommendation 3a	US-guided ablation procedures may be considered in patients with suitable primary papillary microcarcinoma who are unfit for surgery or decline surgery or active surveillance
Recommendation 3b	US-guided ablation procedures may be considered in patients with suitable recurrent papillary thyroid carcinoma who are unfit for surgery or decline surgery or active surveillance
Recommendation 3c	Repeat ablation of a benign nodule can be considered for remnant nodular tissue contributing to unresolved symptomatic or cosmetic concerns

*This is not a comprehensive list of recommendations from the guideline.

RAI: radioactive iodine; RFA: radiofrequency ablation; US: Ultrasound.

American Urological Association

The American Urological Association (AUA; 2017) guideline on renal masses and localized renal cancer affirms that partial nephrectomy should be prioritized for the management of cT1a renal masses when intervention is indicated. Thermal ablation should be considered "as an alternate approach for the management of cT1a renal masses <3 cm in size." The guidelines were updated in 2021 and recommendations are generally consistent with what was published in the 2017 guideline. The 2021 AUA guideline explicitly states that RFA and cryoablation may be offered as options to patients who elect thermal ablation.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

American Thyroid Association

The American Thyroid Association (2015) guideline on the management of thyroid nodules and differentiated thyroid cancer provides recommendations for management. Patients with a benign cytology diagnosis or those very unlikely to be malignant (eg, purely cystic nodule) should undergo surveillance with the frequency determined by the level of suspicion for a missed malignancy. Medical or surgical intervention is considered if the nodules are large (>4 cm), causing compressive or structural symptoms, or if there is clinical concern. Recurrent cystic thyroid nodules with benign cytology should be considered for surgical removal or percutaneous ethanol injection. For differentiated thyroid cancer, "localized treatments with thermal (radiofrequency or cryo-) ablation, ethanol ablation, or chemoembolization may be beneficial in patients with a single or a few metastases and in those with metastases at high risk of local complications."

National Comprehensive Cancer Network

The National Comprehensive Cancer Network (NCCN) guidelines for the treatment of NSCLC (v.3.2023) state: "For medically operable disease, resection is the preferred local treatment modality (other modalities include SABR [stereotactic ablative radiotherapy], thermal ablation such as radiofrequency ablation, and cryotherapy)." For patients who are not amenable to surgery, image-guided thermal ablation therapy (IGTA; includes RFA, microwave ablation, and cryoablation) may be considered. The guidance states "IGTA is an option for the management of NSCLC lesions <3 cm. Ablation for NSCLC lesions >3 cm may be associated with higher rates of local recurrence and complications."

The NCCN guidelines for thyroid carcinoma (v.3.2023) indicate that local therapies such as RFA may be considered for locoregional recurrence of thyroid carcinoma-papillary carcinoma in select patients with limited burden nodal disease. Additionally, local therapies, including RFA, can be considered in those with metastatic disease.

The NCCN guidelines (v.1.2024) for renal cancer indicate that "thermal ablation (eg, cryosurgery, radiofrequency ablation, microwave ablation) is an option for the management of clinical stage T1 renal lesions. Thermal ablation is an option for clinical T1b masses in select patients not eligible for surgery. Biopsy of lesions is recommended to be done prior to or at time of ablation. Ablative techniques may require multiple treatments to achieve the same oncologic outcomes as conventional surgery."

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

The NCCN colon cancer guidelines (v.2.2023) state that “resection is the standard approach for the local treatment of resectable metastatic disease. However, patients with liver or lung oligometastases can also be considered for tumor ablation therapy, particularly in cases that may not be optimal for resection. There is extensive evidence on the use of RFA as a reasonable treatment option for non-surgical candidates and for recurrent disease after hepatectomy with small liver metastases that can be treated with clear margins.”

The NCCN guidelines for head and neck cancers (v.2.2023), breast cancer (v.4.2023), bone cancer (v.1.2024), and pancreatic adenocarcinoma (v.2.2023) do not mention RFA.

National Institute for Health and Care Excellence

The NICE guidance (2004) on osteoid osteoma indicated that "current evidence on the safety and efficacy of computed tomography (CT)-guided thermocoagulation of osteoid osteoma appears adequate to support its use...."

Updated NICE guidance (2010) on renal cancer has indicated that "evidence on the safety and efficacy of percutaneous radiofrequency ablation (RFA) ... in the short and medium term appears adequate to support the use of this procedure provided that patients are followed up in the long term."

The NICE guidance (2010) on RFA for primary and secondary lung cancers has stated: "Current evidence on the efficacy of percutaneous radiofrequency ablation (RFA) ... is adequate in terms of tumor control." The NICE also indicated RFA might "be used in patients with small, early-stage lung cancers or small numbers of lung metastases who are unsuitable for, or prefer not to undergo, surgery. It may also have a place in multi-modality treatment of more advanced primary lung cancers." The guidance warned of serious complications (eg, pneumothorax) among lung cancer patients.

The NICE guidance (2016) on benign thyroid nodules stated: "Current evidence on the safety and efficacy of ultrasound-guided percutaneous radiofrequency ablation ... is adequate to support the use of this procedure...."

Society of Interventional Radiology

The Society of Interventional Radiology (2020) published a position statement on the role of percutaneous ablation in renal cell carcinoma. Their relevant recommendations are as follows:

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

- "In patients with small renal tumors (stage T1a), percutaneous thermal ablation is a safe and effective treatment with fewer complications than nephrectomy and acceptable long-term oncological and survival outcomes. (Level of Evidence: C; Strength of Recommendation: Moderate)"
- "In selected patients with suspected T1a renal cell carcinoma, percutaneous thermal ablation should be offered over active surveillance. (Level of Evidence: C; Strength of Recommendation: Moderate)"
- "In high-risk patients with T1b renal cell carcinoma who are not surgical candidates, percutaneous thermal ablation may be an appropriate treatment option; however, further research in this area is required. (Level of Evidence: D; Strength of Recommendation: Weak)"
- "Radiofrequency ablation, cryoablation, and microwave ablation are all appropriate modalities for thermal ablation, and method of ablation should be left to the discretion of the operating physician. (Level of Evidence: D; Strength of Recommendation: Weak)"

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 2.

Table 2. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT05189821	RFA Treatment for Papillary Thyroid Microcarcinoma Cohort	50	Nov 2026

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

NCT05189808	Radiofrequency Ablation for Indeterminate Bethesda III Thyroid Nodules	50	Aug 2024
NCT03808779	A Multicenter, Randomized and Controlled Trial of Radiofrequency Ablation vs. Conventional Surgery as Treatment of Papillary Thyroid Microcarcinoma (PTMC)	200	Feb 2024
NCT04619472	A Multicenter, Single Group Target Value Clinical Study to Evaluate Safety and Effectiveness of Radiofrequency Ablation System in the Treatment of Peripheral Lung Tumors	126	May 2023
<i>Unpublished</i>			
NCT01051037	Phase II Study Evaluating Safety and Efficacy of Stereotactic Body Radiotherapy and Radiofrequency Ablation for Medically Inoperable and Recurrent Lung Tumors Near Central Airways	17	Dec 2017 (completed)

NCT: national clinical trial.

References

1. Anthem Blue Cross, Clinical UM Guideline. CG-SURG-61, Cryosurgical, Radiofrequency, Microwave or Laser Ablation to Treat Solid Tumors Outside the Liver. 01/03/2024.
2. Jatoi I, Sung H, Jemal A. The Emergence of the Racial Disparity in U.S. Breast-Cancer Mortality. *N Engl J Med.* Jun 23 2022; 386(25): 2349-2352. PMID 35713541
3. Yedjou CG, Sims JN, Miele L, et al. Health and Racial Disparity in Breast Cancer. *Adv Exp Med Biol.* 2019; 1152: 31-49. PMID 31456178
4. National Cancer Institute. SEER Cancer Stat Facts: Kidney and Renal Pelvis Cancer. 2023. <https://seer.cancer.gov/statfacts/html/kidrp.html>.
5. Howard JM, Nandy K, Woldu SL, et al. Demographic Factors Associated With Non-Guideline-Based Treatment of Kidney Cancer in the United States. *JAMA Netw Open.* Jun 01 2021; 4(6): e2112813. PMID 34106265
6. Levy J, Hopkins T, Morris J, et al. Radiofrequency Ablation for the Palliative Treatment of Bone Metastases: Outcomes from the Multicenter OsteoCool Tumor Ablation Post-Market Study

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

- (OPuS One Study) in 100 Patients. *J Vasc Interv Radiol*. Nov 2020; 31(11): 1745-1752. PMID 33129427
7. Goetz MP, Callstrom MR, Charboneau JW, et al. Percutaneous image-guided radiofrequency ablation of painful metastases involving bone: a multicenter study. *J Clin Oncol*. Jan 15 2004; 22(2): 300-6. PMID 14722039
 8. Grönemeyer DH, Schirp S, Gevargez A. Image-guided radiofrequency ablation of spinal tumors: preliminary experience with an expandable array electrode. *Cancer J*. 2002; 8(1): 33-9. PMID 11898806
 9. Kojima H, Tanigawa N, Kariya S, et al. Clinical assessment of percutaneous radiofrequency ablation for painful metastatic bone tumors. *Cardiovasc Intervent Radiol*. 2006; 29(6): 1022-6. PMID 16988875
 10. Tordjman M, Perronne L, Madelin G, et al. CT-guided radiofrequency ablation for osteoid osteomas: a systematic review. *Eur Radiol*. Nov 2020; 30(11): 5952-5963. PMID 32518986
 11. Lanza E, Thouvenin Y, Viala P, et al. Osteoid osteoma treated by percutaneous thermal ablation: when do we fail? A systematic review and guidelines for future reporting. *Cardiovasc Intervent Radiol*. Dec 2014; 37(6): 1530-9. PMID 24337349
 12. Albisinni U, Facchini G, Spinnato P, et al. Spinal osteoid osteoma: efficacy and safety of radiofrequency ablation. *Skeletal Radiol*. Aug 2017; 46(8): 1087-1094. PMID 28497160
 13. Lassalle L, Campagna R, Corcos G, et al. Therapeutic outcome of CT-guided radiofrequency ablation in patients with osteoid osteoma. *Skeletal Radiol*. Jul 2017; 46(7): 949-956. PMID 28429047
 14. Rimondi E, Mavrogenis AF, Rossi G, et al. Radiofrequency ablation for non-spinal osteoid osteomas in 557 patients. *Eur Radiol*. Jan 2012; 22(1): 181-8. PMID 21842430
 15. Sahin C, Oc Y, Ediz N, et al. The safety and the efficacy of computed tomography guided percutaneous radiofrequency ablation of osteoid osteoma. *Acta Orthop Traumatol Turc*. Sep 2019; 53(5): 360-365. PMID 31371131
 16. Knudsen M, Riishede A, Lücke A, et al. Computed tomography-guided radiofrequency ablation is a safe and effective treatment of osteoid osteoma located outside the spine. *Dan Med J*. May 2015; 62(5). PMID 26050823
 17. Rosenthal DI, Hornicek FJ, Torriani M, et al. Osteoid osteoma: percutaneous treatment with radiofrequency energy. *Radiology*. Oct 2003; 229(1): 171-5. PMID 12944597
 18. Yanagisawa T, Mori K, Kawada T, et al. Differential efficacy of ablation therapy versus partial nephrectomy between clinical T1a and T1b renal tumors: A systematic review and meta-analysis. *Urol Oncol*. Jul 2022; 40(7): 315-330. PMID 35562311

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

19. Uhlig J, Strauss A, Rücker G, et al. Partial nephrectomy versus ablative techniques for small renal masses: a systematic review and network meta-analysis. *Eur Radiol.* Mar 2019; 29(3): 1293-1307. PMID 30255245
20. Katsanos K, Mailli L, Krokidis M, et al. Systematic review and meta-analysis of thermal ablation versus surgical nephrectomy for small renal tumours. *Cardiovasc Intervent Radiol.* Apr 2014; 37(2): 427-37. PMID 24482030
21. El Dib R, Touma NJ, Kapoor A. Cryoablation vs radiofrequency ablation for the treatment of renal cell carcinoma: a meta-analysis of case series studies. *BJU Int.* Aug 2012; 110(4): 510-6. PMID 22304329
22. Liu SY, Chu CM, Kong AP, et al. Radiofrequency ablation compared with laparoscopic adrenalectomy for aldosterone-producing adenoma. *Br J Surg.* Oct 2016; 103(11): 1476-86. PMID 27511444
23. Marshall HR, Shakeri S, Hosseiny M, et al. Long-Term Survival after Percutaneous Radiofrequency Ablation of Pathologically Proven Renal Cell Carcinoma in 100 Patients. *J Vasc Interv Radiol.* Jan 2020; 31(1): 15-24. PMID 31767409
24. Andrews JR, Atwell T, Schmit G, et al. Oncologic Outcomes Following Partial Nephrectomy and Percutaneous Ablation for cT1 Renal Masses. *Eur Urol.* Aug 2019; 76(2): 244-251. PMID 31060824
25. Park BK, Gong IH, Kang MY, et al. RFA versus robotic partial nephrectomy for T1a renal cell carcinoma: a propensity score-matched comparison of mid-term outcome. *Eur Radiol.* Jul 2018; 28(7): 2979-2985. PMID 29426988
26. Dai Y, Covarrubias D, Uppot R, et al. Image-Guided Percutaneous Radiofrequency Ablation of Central Renal Cell Carcinoma: Assessment of Clinical Efficacy and Safety in 31 Tumors. *J Vasc Interv Radiol.* Dec 2017; 28(12): 1643-1650. PMID 28673657
27. Dvorak P, Hoffmann P, Brodak M, et al. Percutaneous radiofrequency and microwave ablation in the treatment of renal tumors - 10 years of experience. *Wideochir Inne Tech Maloinwazyjne.* Dec 2017; 12(4): 394-402. PMID 29362655
28. Pantelidou M, Challacombe B, McGrath A, et al. Percutaneous Radiofrequency Ablation Versus Robotic-Assisted Partial Nephrectomy for the Treatment of Small Renal Cell Carcinoma. *Cardiovasc Intervent Radiol.* Nov 2016; 39(11): 1595-1603. PMID 27435582
29. Iannuccilli JD, Dupuy DE, Beland MD, et al. Effectiveness and safety of computed tomography-guided radiofrequency ablation of renal cancer: a 14-year single institution experience in 203 patients. *Eur Radiol.* Jun 2016; 26(6): 1656-64. PMID 26373755

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

30. Schlijper RC, Grutters JP, Houben R, et al. What to choose as radical local treatment for lung metastases from colo-rectal cancer: surgery or radiofrequency ablation?. *Cancer Treat Rev.* Feb 2014; 40(1): 60-7. PMID 23768754
31. Ratko TA, Vats V, Brock J, et al. *Local Nonsurgical Therapies for Stage I and Symptomatic Obstructive Non- Small-Cell Lung Cancer (Comparative Effectiveness Review No. 112)*. Rockville, MD: Agency for Healthcare Research and Quality; 2013.
32. Bilal H, Mahmood S, Rajashanker B, et al. Is radiofrequency ablation more effective than stereotactic ablative radiotherapy in patients with early stage medically inoperable non-small cell lung cancer?. *Interact Cardiovasc Thorac Surg.* Aug 2012; 15(2): 258-65. PMID 22581864
33. Chan VO, McDermott S, Malone DE, et al. Percutaneous radiofrequency ablation of lung tumors: evaluation of the literature using evidence-based techniques. *J Thorac Imaging.* Feb 2011; 26(1): 18-26. PMID 20829720
34. Huang L, Han Y, Zhao J, et al. Is radiofrequency thermal ablation a safe and effective procedure in the treatment of pulmonary malignancies?. *Eur J Cardiothorac Surg.* Mar 2011; 39(3): 348-51. PMID 20663679
35. Zemlyak A, Moore WH, Bilfinger TV. Comparison of survival after sublobar resections and ablative therapies for stage I non-small cell lung cancer. *J Am Coll Surg.* Jul 2010; 211(1): 68-72. PMID 20610251
36. Lencioni R, Crocetti L, Cioni R, et al. Response to radiofrequency ablation of pulmonary tumours: a prospective, intention-to-treat, multicentre clinical trial (the RAPTURE study). *Lancet Oncol.* Jul 2008; 9(7): 621-8. PMID 18565793
37. Zhu JC, Yan TD, Glenn D, et al. Radiofrequency ablation of lung tumors: feasibility and safety. *Ann Thorac Surg.* Apr 2009; 87(4): 1023-8. PMID 19324122
38. Pennathur A, Abbas G, Gooding WE, et al. Image-guided radiofrequency ablation of lung neoplasm in 100 consecutive patients by a thoracic surgical service. *Ann Thorac Surg.* Nov 2009; 88(5): 1601-6; discussion 1607-8. PMID 19853119
39. Xia LY, Hu QL, Xu WY. Efficacy and Safety of Radiofrequency Ablation for Breast Cancer Smaller Than 2 cm: A Systematic Review and Meta-Analysis. *Front Oncol.* 2021; 11: 651646. PMID 34012918
40. Peek MCL, Ahmed M, Napoli A, et al. Minimally invasive ablative techniques in the treatment of breast cancer: a systematic review and meta-analysis. *Int J Hyperthermia.* Mar 2017; 33(2): 191-202. PMID 27575566
41. Zhao Z, Wu F. Minimally-invasive thermal ablation of early-stage breast cancer: a systemic review. *Eur J Surg Oncol.* Dec 2010; 36(12): 1149-55. PMID 20889281

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

42. Soukup B, Bismohun S, Reefy S, et al. The evolving role of radiofrequency ablation therapy of breast lesions. *Anticancer Res.* Sep 2010; 30(9): 3693-7. PMID 20944155
43. Ito T, Oura S, Nagamine S, et al. Radiofrequency Ablation of Breast Cancer: A Retrospective Study. *Clin Breast Cancer.* Aug 2018; 18(4): e495-e500. PMID 29079443
44. Li P, Xiao-Yin T, Cui D, et al. Evaluation of the safety and efficacy of percutaneous radiofrequency ablation for treating multiple breast fibroadenoma. *J Cancer Res Ther.* Dec 2016; 12(Supplement): C138-C142. PMID 28230006
45. Wilson M, Korourian S, Boneti C, et al. Long-term results of excision followed by radiofrequency ablation as the sole means of local therapy for breast cancer. *Ann Surg Oncol.* Oct 2012; 19(10): 3192-8. PMID 22911363
46. Kinoshita T, Iwamoto E, Tsuda H, et al. Radiofrequency ablation as local therapy for early breast carcinomas. *Breast Cancer.* Jan 2011; 18(1): 10-7. PMID 20072824
47. Imoto S, Wada N, Sakemura N, et al. Feasibility study on radiofrequency ablation followed by partial mastectomy for stage I breast cancer patients. *Breast.* Apr 2009; 18(2): 130-4. PMID 19324550
48. Garbay JR, Mathieu MC, Lamuraglia M, et al. Radiofrequency thermal ablation of breast cancer local recurrence: a phase II clinical trial. *Ann Surg Oncol.* Nov 2008; 15(11): 3222-6. PMID 18709415
49. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid.* Jan 2016; 26(1): 1-133. PMID 26462967
50. Cho SJ, Baek JH, Chung SR, et al. Long-Term Results of Thermal Ablation of Benign Thyroid Nodules: A Systematic Review and Meta-Analysis. *Endocrinol Metab (Seoul).* Jun 2020; 35(2): 339-350. PMID 32615718
51. Chen F, Tian G, Kong D, et al. Radiofrequency ablation for treatment of benign thyroid nodules: A PRISMA-compliant systematic review and meta-analysis of outcomes. *Medicine (Baltimore).* Aug 2016; 95(34): e4659. PMID 27559968
52. Fuller CW, Nguyen SA, Lohia S, et al. Radiofrequency ablation for treatment of benign thyroid nodules: systematic review. *Laryngoscope.* Jan 2014; 124(1): 346-53. PMID 24122763
53. Kim JH, Yoo WS, Park YJ, et al. Efficacy and Safety of Radiofrequency Ablation for Treatment of Locally Recurrent Thyroid Cancers Smaller than 2 cm. *Radiology.* Sep 2015; 276(3): 909-18. PMID 25848897

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

54. Owen RP, Khan SA, Negassa A, et al. Radiofrequency ablation of advanced head and neck cancer. *Arch Otolaryngol Head Neck Surg.* May 2011; 137(5): 493-8. PMID 21576561
55. Brook AL, Gold MM, Miller TS, et al. CT-guided radiofrequency ablation in the palliative treatment of recurrent advanced head and neck malignancies. *J Vasc Interv Radiol.* May 2008; 19(5): 725-35. PMID 18440462
56. Owen RP, Silver CE, Ravikumar TS, et al. Techniques for radiofrequency ablation of head and neck tumors. *Arch Otolaryngol Head Neck Surg.* Jan 2004; 130(1): 52-6. PMID 14732768
57. Rey VE, Labrador R, Falcon M, et al. Transvaginal Radiofrequency Ablation of Myomas: Technique, Outcomes, and Complications. *J Laparoendosc Adv Surg Tech A.* Jan 2019; 29(1): 24-28. PMID 30198831
58. Yin G, Chen M, Yang S, et al. Treatment of uterine myomas by radiofrequency thermal ablation: a 10-year retrospective cohort study. *Reprod Sci.* May 2015; 22(5): 609-14. PMID 25355802
59. Liu B, Mo C, Wang W, et al. Treatment outcomes of percutaneous radiofrequency ablation versus adrenalectomy for adrenal metastases: a retrospective comparative study. *J Endocrinol Invest.* Sep 2020; 43(9): 1249-1257. PMID 32166699
60. Yang MH, Tyan YS, Huang YH, et al. Comparison of radiofrequency ablation versus laparoscopic adrenalectomy for benign aldosterone-producing adenoma. *Radiol Med.* Oct 2016; 121(10): 811-9. PMID 27300650
61. Hasegawa T, Takaki H, Kodama H, et al. Three-year Survival Rate after Radiofrequency Ablation for Surgically Resectable Colorectal Lung Metastases: A Prospective Multicenter Study. *Radiology.* Mar 2020; 294(3): 686-695. PMID 31934829
62. Locklin JK, Mannes A, Berger A, et al. Palliation of soft tissue cancer pain with radiofrequency ablation. *J Support Oncol.* 2004; 2(5): 439-45. PMID 15524075
63. Rosenthal DI. Radiofrequency treatment. *Orthop Clin North Am.* Jul 2006; 37(3): 475-84, viii. PMID 16846772
64. Liapi E, Geschwind JF. Transcatheter and ablative therapeutic approaches for solid malignancies. *J Clin Oncol.* Mar 10 2007; 25(8): 978-86. PMID 17350947
65. Spiliotis JD, Datsis AC, Michalopoulos NV, et al. Radiofrequency ablation combined with palliative surgery may prolong survival of patients with advanced cancer of the pancreas. *Langenbecks Arch Surg.* Jan 2007; 392(1): 55-60. PMID 17089173
66. Zou YP, Li WM, Zheng F, et al. Intraoperative radiofrequency ablation combined with 125 iodine seed implantation for unresectable pancreatic cancer. *World J Gastroenterol.* Oct 28 2010; 16(40): 5104-10. PMID 20976848

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

67. Cantore M, Girelli R, Mambrini A, et al. Combined modality treatment for patients with locally advanced pancreatic adenocarcinoma. *Br J Surg.* Aug 2012; 99(8): 1083-8. PMID 22648697
68. Rombouts SJ, Vogel JA, van Santvoort HC, et al. Systematic review of innovative ablative therapies for the treatment of locally advanced pancreatic cancer. *Br J Surg.* Feb 2015; 102(3): 182-93. PMID 25524417
69. Kameyama S, Murakami H, Masuda H, et al. Minimally invasive magnetic resonance imaging-guided stereotactic radiofrequency thermocoagulation for epileptogenic hypothalamic hamartomas. *Neurosurgery.* Sep 2009; 65(3): 438-49; discussion 449. PMID 19687687
70. Vavra P, Dostalík J, Zacharoulis D, et al. Endoscopic radiofrequency ablation in colorectal cancer: initial clinical results of a new bipolar radiofrequency ablation device. *Dis Colon Rectum.* Feb 2009; 52(2): 355-8. PMID 19279436
71. Mylona S, Karagiannis G, Patsoura S, et al. Palliative treatment of rectal carcinoma recurrence using radiofrequency ablation. *Cardiovasc Intervent Radiol.* Aug 2012; 35(4): 875-82. PMID 22167304
72. Ripley RT, Gajdos C, Reppert AE, et al. Sequential radiofrequency ablation and surgical debulking for unresectable colorectal carcinoma: thermo-surgical ablation. *J Surg Oncol.* Feb 2013; 107(2): 144-7. PMID 22927225
73. Howington JA, Blum MG, Chang AC, et al. Treatment of stage I and II non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest.* May 2013; 143(5 Suppl): e278S-e313S. PMID 23649443
74. Donington J, Ferguson M, Mazzone P, et al. American College of Chest Physicians and Society of Thoracic Surgeons consensus statement for evaluation and management for high-risk patients with stage I non-small cell lung cancer. *Chest.* Dec 2012; 142(6): 1620-1635. PMID 23208335
75. Orloff LA, Noel JE, Stack BC, et al. Radiofrequency ablation and related ultrasound-guided ablation technologies for treatment of benign and malignant thyroid disease: An international multidisciplinary consensus statement of the American Head and Neck Society Endocrine Surgery Section with the Asia Pacific Society of Thyroid Surgery, Associazione Medici Endocrinologi, British Association of Endocrine and Thyroid Surgeons, European Thyroid Association, Italian Society of Endocrine Surgery Units, Korean Society of Thyroid Radiology, Latin American Thyroid Society, and Thyroid Nodules Therapies Association. *Head Neck.* Mar 2022; 44(3): 633-660. PMID 34939714
76. Campbell S, Uzzo RG, Allaf ME, et al. Renal Mass and Localized Renal Cancer: AUA Guideline. *J Urol.* Sep 2017; 198(3): 520-529. PMID 28479239

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

77. Campbell SC, Clark PE, Chang SS, et al. Renal Mass and Localized Renal Cancer: Evaluation, Management, and Follow-Up: AUA Guideline: Part I. J Urol. Aug 2021; 206(2): 199-208. PMID 34115547
78. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Non-small cell lung cancer. Version 3.2023. Updated April 13, 2023. https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf.
79. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Thyroid Carcinoma. Version 3.2023. Updated July 27, 2023. https://www.nccn.org/professionals/physician_gls/pdf/thyroid.pdf.
80. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Kidney Cancer. Version 1.2024. Updated June 21, 2023. https://www.nccn.org/professionals/physician_gls/pdf/kidney.pdf.
81. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Colon Cancer. Version 2.2023. Updated April 25, 2023. https://www.nccn.org/professionals/physician_gls/pdf/colon.pdf.
82. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Head and Neck Cancers. Version 2.2023. Updated May 15, 2023. https://www.nccn.org/professionals/physician_gls/pdf/head-and-neck.pdf.
83. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Breast Cancer. Version 4.2023. Updated March 23, 2023. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf.
84. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Bone Cancer. Version 3.2023. Updated April 4, 2023. https://www.nccn.org/professionals/physician_gls/pdf/bone.pdf.
85. National Comprehensive Cancer Network. NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines): Pancreatic Adenocarcinoma. Version 2.2023. Updated June 19, 2023. https://www.nccn.org/professionals/physician_gls/pdf/pancreatic.pdf.
86. National Institute for Health and Care Excellence (NICE). Computed tomography-guided thermocoagulation of osteoid osteoma [IPG53]. 2004; <https://www.nice.org.uk/guidance/ipg53>.
87. National Institute for Health and Care Excellence (NICE). Percutaneous radiofrequency ablation of renal cancer [IPG353]. 2010; <https://www.nice.org.uk/guidance/ipg353>.
88. National Institute for Health and Care Excellence (NICE). Percutaneous radiofrequency ablation for primary and secondary lung cancers [IPG372]. 2010; <https://www.nice.org.uk/guidance/ipg372>.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

- 89. National Institute for Health and Care Excellence (NICE). Ultrasound-guided percutaneous radiofrequency ablation for benign thyroid nodules [IPG562]. 2016; <https://www.nice.org.uk/guidance/IPG562>.
- 90. Morris CS, Baerlocher MO, Dariushnia SR, et al. Society of Interventional Radiology Position Statement on the Role of Percutaneous Ablation in Renal Cell Carcinoma: Endorsed by the Canadian Association for Interventional Radiology and the Society of Interventional Oncology. J Vasc Interv Radiol. Feb 2020; 31(2): 189-194.e3. PMID 31917025

Policy History

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

- 07/13/2005 Medical Director review
- 07/19/2005 Medical Policy Committee review
- 08/24/2005 Managed Care Advisory Council approval
- 07/07/2006 Format revision, including addition of FDA and or other governmental regulatory approval and rationale/source. Coverage eligibility unchanged.
- 08/08/2006 Medical Director review
- 08/09/2006 Medical Policy Committee approval.
- 07/10/2007 Medical Director review
- 07/18/2007 Medical Policy Committee approval. Policy Statement added for osteoid osteoma and rationale/source updated.
- 07/02/2008 Medical Director review
- 07/16/2008 Medical Policy Committee approval. Coverage eligibility unchanged.
- 07/02/2009 Medical Director review
- 07/22/2009 Medical Policy Committee approval. Coverage statement added to indicate that use of radiofrequency ablation as treatment of osteoid osteomas that cannot be managed successfully with medical treatment may be considered eligible for coverage. Coverage statement added that treatment of localized renal cell carcinoma no more than 4cm in size may be considered eligible for coverage when specific criteria are met. Added that radiofrequency ablation for the treatment of osteoid osteomas that can be managed with medical treatment is considered to be investigational.
- 08/05/2010 Medical Policy Committee review

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

08/18/2010	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
08/04/2011	Medical Policy Committee review
08/17/2011	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
08/02/2012	Medical Policy Committee review
08/15/2012	Medical Policy Implementation Committee approval. Added options for primary and metastatic pulmonary tumors to be eligible for coverage with criteria.
12/12/2013	Medical Policy Committee review
12/18/2013	Medical Policy Implementation Committee approval. Reformatted investigational statement and added thyroid as investigational.
06/25/2015	Medical Policy Committee review
07/15/2015	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
06/30/2016	Medical Policy Committee review
07/20/2016	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
01/01/2017	Coding update: Removing ICD-9 Diagnosis codes
07/06/2017	Medical Policy Committee review
07/19/2017	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
01/01/2018	Coding update
07/05/2018	Medical Policy Committee review
07/11/2018	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
08/31/2018	Coding update
07/03/2019	Medical Policy Committee review
07/18/2019	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
11/13/2019	Coding update
01/14/2020	Coding update
07/02/2020	Medical Policy Committee review
07/08/2020	Medical Policy Implementation Committee approval. Added an eligible for coverage statement for RFA to treat osteoid osteomas that cannot be managed

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

successfully with medical treatment. Added a reference to the Policy Guidelines after the eligible for coverage criteria for treatment of malignant nonpulmonary tumor(s) metastatic to the lung.

07/01/2021 Medical Policy Committee review

07/14/2021 Medical Policy Implementation Committee approval. Added pancreas to the list for radiofrequency ablation of all other tumors to be investigational. Coverage eligibility unchanged.

07/07/2022 Medical Policy Committee review

07/13/2022 Medical Policy Implementation Committee approval. Coverage eligibility unchanged.

07/06/2023 Medical Policy Committee review

07/12/2023 Medical Policy Implementation Committee approval. Revised the coverage section.

07/02/2024 Medical Policy Committee review

07/10/2024 Medical Policy Implementation Committee approval. Removed thyroid carcinoma as eligible for coverage. Removed biopsy proven requirement for non-small cell lung cancer. Deleted first criteria bullet for metastatic malignant lung tumor(s) for surgical or radiation treatment with curative intent. Metastatic malignant lung tumor(s) criteria with 1st bullet added regarding preserving lung function and surgical candidacy, and criteria for extrapulmonary metastatic disease and for number of tumors per lung were reworded. Added breast and thyroid tumors to the investigational statement.

Next Scheduled Review Date: 07/2025

Coding

The five character codes included in the Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2023 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines is with Blue Cross and Blue Shield of Louisiana and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

attributable or related to any use, nonuse or interpretation of information contained in Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Blue Cross Blue Shield of Louisiana Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	20982, 32998, 50542, 50592, 60699, 76940, 77013, 77022 Delete code effective 09/30/2024: 20983
HCPCS	No codes
ICD-10 Diagnosis	All related Diagnoses

*Investigational – A medical treatment, procedure, drug, device, or biological product is Investigational if the effectiveness has not been clearly tested and it has not been incorporated into standard medical practice. Any determination we make that a medical treatment, procedure, drug, device, or biological product is Investigational will be based on a consideration of the following:

- A. Whether the medical treatment, procedure, drug, device, or biological product can be lawfully marketed without approval of the U.S. Food and Drug Administration (FDA) and whether such approval has been granted at the time the medical treatment, procedure, drug, device, or biological product is sought to be furnished; or
- B. Whether the medical treatment, procedure, drug, device, or biological product requires further studies or clinical trials to determine its maximum tolerated dose, toxicity, safety, effectiveness, or effectiveness as compared with the standard means of treatment or diagnosis, must improve health outcomes, according to the consensus of opinion among experts as shown by reliable evidence, including:

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Louisiana

Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

1. Consultation with technology evaluation center(s);
2. Credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community; or
3. Reference to federal regulations.

****Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.



Radiofrequency Ablation of Miscellaneous Solid Tumors Excluding Liver Tumors

Policy # 00175

Original Effective Date: 08/24/2005

Current Effective Date: 10/01/2024

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

©2024 Blue Cross and Blue Shield of Louisiana

Blue Cross and Blue Shield of Louisiana is an independent licensee of the Blue Cross and Blue Shield Association and incorporated as Louisiana Health Service & Indemnity Company.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, or otherwise, without permission from Blue Cross and Blue Shield of Louisiana.