

Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

Applies to all products administered or underwritten by Blue Cross and Blue Shield of Louisiana and its subsidiary, HMO Louisiana, Inc. (collectively referred to as the "Company"), unless otherwise provided in the applicable contract. Medical technology is constantly evolving, and we reserve the right to review and update Medical Policy periodically.

When Services May Be Eligible for Coverage

Coverage for eligible medical treatments or procedures, drugs, devices or biological products may be provided only if:

- *Benefits are available in the member's contract/certificate, and*
- *Medical necessity criteria and guidelines are met.*

Based on review of available data, the Company may consider the use of an implanted hydrogel spacer between the prostate and rectum when primary definitive radiation therapy will be used to treat prostate cancer using any form of external beam radiation therapy (3D conformal, IMRT, SBRT) to be **eligible for coverage:****

When Services Are Considered Not Medically Necessary

Based on review of available data, the Company considers the use of an implanted hydrogel spacer for all other indications to be **not medically necessary.****

Background/Overview

Radiation Oncology Considerations

Because the anterior wall of the rectum abuts the posterior prostate, radiotherapy for prostate cancer exposes that portion of the rectum to the full dose of radiation delivered to the prostate, which poses the risk of rectal bleeding for months to years after treatment. Modern radiation planning techniques, such as intensity modulated radiation therapy (IMRT), allow significantly higher doses of radiation to be safely delivered to the prostate while maintaining an acceptable risk of late rectal complications by limiting the portion of the rectum treated to full dose. In recent years, attempts to reduce rectal toxicity have focused on increasing the physical distance between the prostate and rectum by injection of a biodegradable hydrogel to push the rectum away from the high dose region to allow additional dose sparing.

The use of an implanted hydrogel spacer between the prostate and rectum has been studied as a way to minimize rectal symptoms during and after definitive radiotherapy for adenocarcinoma of the prostate. A pilot study authored by Song et al. documented the ability to increase the space between the prostate and rectum to an average of 7.5 mm. The additional space resulted in significant reductions in rectal dose across the range of 10 Gy to 75 Gy. No clinical outcomes were reported. Favorable early gastrointestinal (GI) and genitourinary (GU) toxicity profiles were reported in a phase II study by Uhl,

Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

but there was no control group for comparison.

External Beam Radiation Therapy

The only randomized controlled trial (RCT) of hydrogel spacer placement was reported by Mariados. It randomly assigned patients 2:1 for either spacer placement or placebo. Study participants had stage T1 or T2 stage prostate cancer without extracapsular extension. A total of 149 patients had the spacer placed prior to radiotherapy and were compared to 73 patients treated without spacer injection. Both groups were treated with image-guided IMRT to a dose of 79.2 Gy in 44 fractions.

The initial report was published in 2015 and showed no significant reduction in rectal adverse events in the first 6 months (34.2% with spacer vs 31.5% without, $P=.7$). Significant reduction in late (3-15 month) rectal toxicity was associated with spacer placement, with 2% (3 patients) and 7% (5 patients) experiencing grade 1 or greater GI symptoms in the hydrogel and control arms ($P=.044$), respectively. Urinary toxicity was not significantly different between the groups.

Hamstra et al. subsequently reported 36-month results of a subset of the original trial participants. They reported a 0% grade 2 or higher rectal toxicity with spacer use versus a 5.7% rate without the spacer ($P=.012$). They also noted a significant reduction in grade 1 urinary incontinence favoring spacer placement (15% vs 4%, $P=.046$). A subsequent analysis reported an improvement in sexual function with the spacer, but this did not meet statistical significance.

There is a strong secular trend toward the use of shorter courses of external beam radiation therapy to treat low-risk and intermediate-risk prostate cancer. Multiple randomized controlled trials (RCT) of shorter course radiation, also called hypofractionated radiation, have shown equivalent cure rates to conventionally fractionated radiation but with a higher incidence acute rectal toxicity. Given the higher GI toxicity of this regimen, the use of a hydrogel spacer would be most advantageous in this cohort of patients and has become standard of care in this setting.

Stereotactic Body Radiation Therapy

Stereotactic body radiation therapy (SBRT), also termed ultra hypofractionated radiation therapy is an alternative radiation modality to treat low-risk and intermediate risk prostate cancer. Treatment is given in 5 or fewer daily sessions or fractions. Fried et al. reported on the use of a perirectal hydrogel spacer in association with SBRT. The retrospective report demonstrated significant improvement in rectal and penile bulb dosimetry with the use of the spacer in 66 patients compared to 28 patients who had not undergone spacer placement.

A much larger study by Zelefsky and colleagues examined outcomes in 551 patients with low-risk and intermediate-risk prostate cancer treated with SBRT. The treatment consisted of 37.5-40 Gy in 5 fractions directed to the prostate and seminal vesicles. About half of the patients (269/551) received a rectal spacer as this became a standard part of the group's treatment protocol in November 2016. The use of a spacer was associated with a significant reduction in any late GI toxicity (1% with spacer vs 6% without, $P=.010$). Spacer placement also significantly reduced late GU toxicity (15% for spacer vs 32%



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

without, $P < .001$).

Brachytherapy

The use of a hydrogel spacer in the setting of low dose rate (LDR) brachytherapy has been reported by Khan et al. Forty patients who underwent perirectal hydrogel injection were compared to 40 patients who had not undergone spacer placement. Some of the patients also received external beam radiation. There was a reduction in rectal toxicity at 1 month, but no difference in toxicity at either one or 2-year follow-up. This finding was similar to a previous report by Taggar et al. comparing toxicity in 74 patients with spacer placement prior to Pd-103 LDR brachytherapy to a similar cohort without spacers. Similarly, a report by Lin et al. examining non-randomized outcomes of hydrogel spacer use prior to LDR brachytherapy showed reduced rates of grade 1 toxicity but no significant difference in grade 2 or 3 toxicities. Despite improvements in rectal dosimetry, there was no significant improvement in acute rectal toxicity. Further studies are needed to define the role of hydrogel spacer placement, if any, in the setting of brachytherapy.

Systematic Reviews

A systematic review of the use of a hydrogel spacer to reduce toxicity during and after radiotherapy for prostate cancer was recently published by Armstrong et al. This review is more extensive than previous reviews by Miller et al. and the Canadian Agency for Drugs and Technologies in Health (CADTH). In addition to the RCT described above, they reviewed 18 additional spacer studies looking at several radiotherapy techniques. Seven of the 18 studies evaluated hydrogel use with conventionally fractionated IMRT. Two studies examined outcomes when used with SBRT, and one looked at spacer use with proton therapy. Most of the other studies included patients treated with combinations of external beam radiation and brachytherapy.

A recent Cochrane review of interventions to reduce acute and late adverse GI effects of pelvic radiotherapy concluded that “low-certainty evidence on balloon and hydrogel spacers suggests that these interventions for prostate cancer RT may make little or no difference to GI outcomes.”

Toxicity and Risk

A recent commentary published in *Lancet Oncology* urged caution in the widespread use of the hydrogel spacer given the small expected benefit and the rising number of reported adverse events associated with the procedure. Despite excellent safety in the small trial, there are a growing number of reports of significant adverse events in real-world use. By examining the FDA Manufacturer and User Facility Device (MAUDE) database, the authors identified 85 reported events. The majority of these could be converted into graded toxicities using Common Terminology Criteria for Adverse Events. Approximately 70% of the events were graded 3, 4, or 5, with about 24% falling into the grade 4 category, including colostomy, anaphylactic events, rectal wall injection, and pulmonary embolism. There was one death. They concluded that critical reflection and careful consideration of the need, toxicity, and benefits of perirectal hydrogel spacer placement should precede any recommendation for its use.



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

FDA or Other Governmental Regulatory Approval

U.S. Food and Drug Administration (FDA)

In October 2014, SpaceOAR™‡ (Augmenix, a subsidiary of Boston Scientific) was cleared by the U.S. Food and Drug Administration (FDA) through the De Novo process (DEN140030). Barrigel Injectable Gel (Palette Life Sciences) was approved by the FDA via the premarket approval process in March 2022 (K220641; FDA product code: OVB), followed by BioProtect Balloon Implant™‡ System (BioProtect, Ltd) in 2023 (K222972; FDA product code: OVB). The intended and approved use of SpaceOAR System, Barrigel, and BioProtect Balloon Implant is to temporarily position the anterior rectal wall away from the prostate during radiotherapy for prostate cancer and in creating this space it is the intent of these hydrogel spacers to reduce the radiation dose delivered to the anterior rectum.

DuraSeal®‡ Exact (Integra) was approved by the FDA through the premarket approval process as a spine and cranial sealant (dura mater) and has been used off-label as a perirectal spacer.

Rationale/Source

This medical policy was developed through consideration of peer-reviewed medical literature generally recognized by the relevant medical community, U.S. Food and Drug Administration approval status, nationally accepted standards of medical practice and accepted standards of medical practice in this community, technology evaluation centers, reference to regulations, other plan medical policies, and accredited national guidelines.

Supplemental Information

Practice Guidelines and Position Statements

Guidelines or position statements will be considered for inclusion in ‘Supplemental Information’ if they were issued by, or jointly by, a US professional society, an international society with US representation, or National Institute for Health and Care Excellence (NICE). Priority will be given to guidelines that are informed by a systematic review, include strength of evidence ratings, and include a description of management of conflict of interest.

American College of Radiology

American College of Radiology appropriateness criteria, last reviewed in 2016, for dose-volume constraints for the rectum with external beam radiotherapy are described in Table 1.

Table 1. Dose Constraints for the Rectum With External Beam Radiotherapy

EBRT Dose-Volume	Dose	<15%	<25%	<35%	<50%
Conventional Fractionation	1.8 Gy X 44 fractions (79.2 Gy total)	V75	V70	V65	V60



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

Hypofractionation	2.5 Gy X 25 fractions (70 Gy total)	V74	V69	V64	V59
-------------------	-------------------------------------	-----	-----	-----	-----

EBRT: External beam radiotherapy; Gy: gray.
V100 = volume of structure (X%) receiving 100% of the dose

American Society of Clinical Oncology, the American Urological Association, and the American Society for Radiation Oncology

In 2018, the American Society of Clinical Oncology, the American Urological Association, and the American Society for Radiation Oncology published a joint guideline on hypofractionated radiation therapy for localized prostate cancer. The guideline recommends that men be counseled about the small increased risk of acute gastrointestinal (GI) toxicity with hypofractionation. "Moderately fractionated EBRT [external beam radiotherapy] has a similar risk of acute and late genitourinary and late GI toxicity compared with conventionally fractionated EBRT. However, physicians should discuss the limited follow-up beyond 5 years for most existing RCTs [randomized controlled trials] evaluating moderate hypofractionation." This was a strong recommendation based on high-quality evidence and 100% consensus. Additionally, the guideline mentions that prostate-rectal spacers can be used to allow rectal dose sparing.

National Comprehensive Cancer Network

The National Comprehensive Cancer Network guideline for prostate cancer (v4.2024) provides the following recommendation in principles of radiation therapy (PROS-F), "Overall, the panel believes that biocompatible and biodegradable perirectal spacer materials may be implanted between the prostate and rectum in patients undergoing external radiotherapy with organ-confined prostate cancer in order to displace the rectum from high radiation dose regions."

National Institute for Health and Care Excellence

In 2023, NICE updated their guidance on the biodegradable spacer. The NICE recommendations state that: "Evidence on the safety and efficacy of biodegradable spacer insertion to reduce rectal toxicity during radiotherapy for prostate cancer is limited in quality. Therefore, this procedure should only be used with special arrangements for clinical governance, consent, and audit or research."

U.S. Preventive Services Task Force Recommendations

Not applicable.

Medicare National Coverage

There is no national coverage determination. In the absence of a national coverage determination, coverage decisions are left to the discretion of local Medicare carriers.

Ongoing and Unpublished Clinical Trials

Some currently unpublished trials that might influence this review are listed in Table 2.



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

Table 2. Summary of Key Trials

NCT No.	Trial Name	Planned Enrollment	Completion Date
<i>Ongoing</i>			
NCT04905069 ^a	Effectiveness of the SpaceOAR Vue System in Subjects With Prostate Cancer Being Treated With Stereotactic Body Radiotherapy	500	Apr 2030
NCT05597852	Feasibility of Integrating Rectal Hydrogel Spacer for Salvage Treatment Using Stereotactic Ablative Body Radiotherapy for Locally Recurrent Prostate Cancer	10	Nov 2027
NCT05650021	Radiopaque Hydrogel Rectal Spacer for Prostate Cancer Radiation Image Guidance	30	Sep 2025
NCT05354440 ^a	Long-Term Prospective Post Marketing Clinical Follow Up for Evaluation of the BioProtect Balloon Implant System	80	Jan 2026
<i>Unpublished</i>			
NCT05354427 ^a	Evaluation of Commercially Available Implantable Spacers, in Prostate Cancer Patients Undergoing Radiotherapy	175	Jan 2022
NCT00462124 ^a	One-arm, Multi-center, Prospective Study to Assess the Safety and Efficacy of BioProtect Biodegradable Implantable Balloon in Prostate Cancer Subjects Undergoing Radiotherapy	7	May 2009

NCT: national clinical trial.

^a Denotes industry-sponsored or cosponsored trial.

References

1. Carelon Medical Benefits Management Guidelines Radiation Oncology. Appropriate Use Criteria: Brachytherapy, Intensity Modulated Radiation Therapy, Stereotactic body Radiation Therapy, and Stereotactic Radiosurgery. Effective 03/23/2025.
2. Afkhami Ardekani M, Ghaffari H, Navaser M, et al. Effectiveness of rectal displacement devices in managing prostate motion: a systematic review. *Strahlenther Onkol.* 2021;197(2):97-115.
3. Aminsharifi A, Kotamarti S, Silver D, et al. Major complications and adverse events related to the injection of the SpaceOAR hydrogel system before radiotherapy for prostate cancer: review of the manufacturer and user facility device experience database. *J Endourol.* 2019;33(10):868-71.



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

4. Armstrong N, Bahl A, Pinkawa M, et al. SpaceOAR hydrogel spacer for reducing radiation toxicity during radiotherapy for prostate cancer. a systematic review. *Urology*. 2021;156:e74-e85.
5. Babayan RK, Steinberg ML, Miller LE. Re: Aminsharifi et al., Major complications and adverse events related to the injection of the SpaceOAR hydrogel system before radiotherapy for prostate cancer: review of the manufacturer and user facility device experience database (From: Aminsharifi A, Kotamarti S, Silver D, et al., *J Endourol* 2019;33:868-871; DOI: 10.1089/end.2019.0431). *J Endourol*. 2020;34(2):240-1.
6. Carvalho IT, Baccaglini W, Claros OR, et al. Genitourinary and gastrointestinal toxicity among patients with localized prostate cancer treated with conventional versus moderately hypofractionated radiation therapy: systematic review and meta-analysis. *Acta Oncol*. 2018;57(8):1003-10.
7. Chao Y, MacDougall D, de Nanassy A, et al. CADTH rapid response report: summary with critical appraisal. Hydrogel spacers for patients with prostate cancer: a review of clinical effectiveness and cost-effectiveness. 2019. [36 p.]. Available from: <https://www.cadth.ca/hydrogel-spacers-patients-prostate-cancer-review-clinical-effectiveness-and-cost-effectiveness>.
8. Dearnaley D, Syndikus I, Mossop H, et al. Conventional versus hypofractionated high-dose intensity-modulated radiotherapy for prostate cancer: 5-year outcomes of the randomised, non-inferiority, phase 3 CHHiP trial. *Lancet Oncol*. 2016;17(8):1047-60.
9. Fransson P, Nilsson P, Gunnlaugsson A, et al. Ultra-hypofractionated versus conventionally fractionated radiotherapy for prostate cancer (HYPO-RT-PC): patient-reported quality-of-life outcomes of a randomised, controlled, non-inferiority, phase 3 trial. *Lancet Oncol*. 2021;22(2):235-45.
10. Fried DB, Dubose RS, Johnson K, et al. Dosimetry for organs at risk with and without use of perirectal hydrogel spacer in prostate cancer patients treated with SBRT. *International Journal of Radiation Oncology Biology Physics*. 2017;99 (2 Supplement):abstract 2559.
11. Hall WA, Tree AC, Dearnaley D, et al. Considering benefit and risk before routinely recommending SpaceOAR. *Lancet Oncol*. 2021;22(1):11-3.
12. Hamstra DA, Mariados N, Sylvester J, et al. Sexual quality of life following prostate intensity modulated radiation therapy (IMRT) with a rectal/prostate spacer: secondary analysis of a phase 3 trial. *Pract Radiat Oncol*. 2018;8(1):e7-e15.
13. Hamstra DA, Mariados N, Sylvester J, et al. Continued benefit to rectal separation for prostate radiation therapy: final results of a phase III trial. *Int J Radiat Oncol Biol Phys*. 2017;97(5):976-85.
14. Hickey BE, James ML, Daly T, et al. Hypofractionation for clinically localized prostate cancer (review). *Cochrane Database Syst Rev*. 2019(9):CD011462.
15. Hoffman KE, Voong KR, Levy LB, et al. Randomized trial of hypofractionated, dose-escalated, intensity-modulated radiation therapy (IMRT) versus conventionally fractionated IMRT for localized prostate cancer. *J Clin Oncol*. 2018;36(29):2943-9.
16. Hrobjartsson A, Thomsen AS, Emanuelsson F, et al. Observer bias in randomized clinical trials with measurement scale outcomes: a systematic review of trials with both blinded and nonblinded assessors. *CMAJ*. 2013;185(4):E201-11.
17. Incrocci L, Wortel RC, Alemayehu WG, et al. Hypofractionated versus conventionally fractionated radiotherapy for patients with localised prostate cancer (HYPRO): final efficacy results from a randomised, multicentre, open-label, phase 3 trial. *Lancet Oncol*. 2016;17(8):1061-9.



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

18. Kahn J, Dahman B, McLaughlin C, et al. Rectal spacing, prostate coverage, and periprocedural outcomes after hydrogel spacer injection during low-dose-rate brachytherapy implantation. *Brachytherapy*. 2020;19(2):228-33.
19. King CR, Kamrava M, Wang PC, et al. In regard to Mariados et al. *Int J Radiat Oncol Biol Phys*. 2015;1(4):936-7.
20. Lawrie TA, Green JT, Beresford M, et al. Interventions to reduce acute and late adverse gastrointestinal effects of pelvic radiotherapy for primary pelvic cancers. *Cochrane Database Syst Rev*. 2018(1):article number CD012529.
21. Lewis SC, Warlow CP. How to spot bias and other potential problems in randomised controlled trials. *J Neurol Neurosurg Psychiatry*. 2004;75(2):181-7.
22. Lin YH, Loon W, Tacey M, et al. Impact of hydrogel and hyaluronic acid rectal spacer on rectal dosimetry and toxicity in low-dose-rate prostate brachytherapy: a multi-institutional analysis of patients' outcomes. *J Contemp Brachytherapy*. 2021;13(6):605-14.
23. Mariados N, Hamstra DA. In reply to King et al. *Int J Radiat Oncol Biol Phys*. 2015;1(4):937-9.
24. Mariados N, Sylvester J, Shah D, et al. Hydrogel spacer prospective multicenter randomized controlled pivotal trial: dosimetric and clinical effects of perirectal spacer application in men undergoing prostate image guided intensity modulated radiation therapy. *Int J Radiat Oncol Biol Phys*. 2015;92(5):971-7.
25. Miller LE, Efsthathiou JA, Bhattacharyya SK, et al. Association of the placement of a perirectal hydrogel spacer with the clinical outcomes of men receiving radiotherapy for prostate cancer: a systematic review and meta-analysis. *JAMA Netw Open*. 2020;3(6):e208221.
26. Morgan SC, Hoffman K, Loblaw DA, et al. Hypofractionated radiation therapy for localized prostate cancer: an ASTRO, ASCO, and AUA evidence-based guideline [complete unabridged version of the guideline; supplementary material]. 2018. [52 p.]. Available from: <https://doi.org/10.1016/j.prro.2018.08.002>.
27. Morgan SC, Hoffman K, Loblaw DA, et al. Hypofractionated radiation therapy for localized prostate cancer: executive summary of an ASTRO, ASCO and AUA evidence-based guideline. *J Urol*. 2019;201(3):528-34.
28. Morgan SC, Morton GC, Berlin A, et al. Current topics in radiotherapy for genitourinary cancers: consensus statements of the Genitourinary Radiation Oncologists of Canada. *Can Urol Assoc J*. 2020;14(11):E588-E93.
29. Nossiter J, Sujenthiran A, Cowling TE, et al. Patient-reported functional outcomes after hypofractionated or conventionally fractionated radiation for prostate cancer: a national cohort study in England. *J Clin Oncol*. 2020;38(7):744-52.
30. Ogita M, Yamashita H, Nozawa Y, et al. Phase II study of stereotactic body radiotherapy with hydrogel spacer for prostate cancer: acute toxicity and propensity score-matched comparison. *Radiat Oncol*. 2021;16(1):107.
31. Payne HA, Pinkawa M, Peedell C, et al. SpaceOAR hydrogel spacer injection prior to stereotactic body radiation therapy for men with localized prostate cancer: A systematic review. *Medicine (Baltimore)*. 2021;100(49):e28111.
32. Quinn TJ, Daignault-Newton S, Bosch W, et al. Who benefits from a prostate rectal spacer? secondary analysis of a phase III trial. *Pract Radiat Oncol*. 2020;10(3):186-94.



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

33. Song DY, Herfarth KK, Uhl M, et al. A multi-institutional clinical trial of rectal dose reduction via injected polyethylene-glycol hydrogel during intensity modulated radiation therapy for prostate cancer: analysis of dosimetric outcomes. *Int J Radiat Oncol Biol Phys.* 2013;87(1):81-7.
34. Taggar AS, Charas T, Cohen GN, et al. Placement of an absorbable rectal hydrogel spacer in patients undergoing low-dose-rate brachytherapy with palladium-103. *Brachytherapy.* 2018;17(2):251-8.
35. Uhl M, Herfarth K, Eble MJ, et al. Absorbable hydrogel spacer use in men undergoing prostate cancer radiotherapy: 12 month toxicity and proctoscopy results of a prospective multicenter phase II trial. *Radiat Oncol.* 2014;9:96.
36. Widmark A, Gunnlaugsson A, Beckman L, et al. Ultra-hypofractionated versus conventionally fractionated radiotherapy for prostate cancer: 5-year outcomes of the HYPO-RT-PC randomised, non-inferiority, phase 3 trial. *Lancet.* 2019;394(10196):385-95.
37. Yin Z, You J, Wang Y, et al. Moderate hypofractionated radiotherapy vs conventional fractionated radiotherapy in localized prostate cancer: a systemic review and meta-analysis from Phase III randomized trials. *Onco Targets Ther.* 2019;12:1259-68.
38. Zelefsky MJ, Pinitpatcharalert A, Kollmeier M, et al. Early tolerance and tumor control outcomes with high-dose ultrahypofractionated radiation therapy for prostate cancer. *Eur Urol Oncol.* 2019;3(6):748-55.

Policy History

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

04/04/2019	Medical Policy Committee review
04/24/2019	Medical Policy Implementation Committee approval. New policy.
04/02/2020	Medical Policy Committee review
04/08/2020	Medical Policy Implementation Committee approval. Coverage eligibility unchanged.
04/01/2021	Medical Policy Committee review
04/14/2021	Medical Policy Implementation Committee approval. Coverage changed from investigational to not medically necessary.
11/10/2021	Medical Policy Implementation Committee approval. Coverage changed from Not Medically necessary to eligible with criteria.
11/03/2022	Medical Policy Committee review
11/09/2022	Medical Policy Implementation Committee approval. No change to coverage.
01/05/2023	Medical Policy Committee review
01/11/2023	Medical Policy Implementation Committee approval. No change to coverage.
01/04/2024	Medical Policy Committee review
01/10/2024	Medical Policy Implementation Committee approval. No change to coverage.
01/02/2025	Medical Policy Committee review
01/08/2025	Medical Policy Implementation Committee approval. Removed “low risk or intermediate risk” prostate cancer using “EITHER of the following techniques”. And added “any form of external beam radiation therapy (3D conformal, IMRT, SBRT)” to be eligible for coverage. Also removed patient selection criteria and the policy guidelines sections from policy.



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

Next Scheduled Review Date: 01/2026

Coding

The five character codes included in the Louisiana Blue Medical Policy Coverage Guidelines are obtained from Current Procedural Terminology (CPT®)‡, copyright 2024 by the American Medical Association (AMA). CPT is developed by the AMA as a listing of descriptive terms and five character identifying codes and modifiers for reporting medical services and procedures performed by physician.

The responsibility for the content of Louisiana Blue Medical Policy Coverage Guidelines is with Louisiana Blue and no endorsement by the AMA is intended or should be implied. The AMA disclaims responsibility for any consequences or liability attributable or related to any use, nonuse or interpretation of information contained in Louisiana Blue Medical Policy Coverage Guidelines. Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein. Any use of CPT outside of Louisiana Blue Medical Policy Coverage Guidelines should refer to the most current Current Procedural Terminology which contains the complete and most current listing of CPT codes and descriptive terms. Applicable FARS/DFARS apply.

CPT is a registered trademark of the American Medical Association.

Codes used to identify services associated with this policy may include (but may not be limited to) the following:

Code Type	Code
CPT	55874
HCPCS	No codes
ICD-10 Diagnosis	All related diagnoses

****Medically Necessary (or “Medical Necessity”)** - Health care services, treatment, procedures, equipment, drugs, devices, items or supplies that a Provider, exercising prudent clinical judgment, would provide to a patient for the purpose of preventing, evaluating, diagnosing or treating an illness, injury, disease or its symptoms, and that are:

- A. In accordance with nationally accepted standards of medical practice;
- B. Clinically appropriate, in terms of type, frequency, extent, level of care, site and duration, and considered effective for the patient's illness, injury or disease; and
- C. Not primarily for the personal comfort or convenience of the patient, physician or other health care provider, and not more costly than an alternative service or sequence of services at least as likely to produce equivalent therapeutic or diagnostic results as to the diagnosis or treatment of that patient's illness, injury or disease.



Hydrogel Spacer use During Radiotherapy for Prostate Cancer

Policy # 00662

Original Effective Date: 07/01/2019

Current Effective Date: 03/23/2025

For these purposes, “nationally accepted standards of medical practice” means standards that are based on credible scientific evidence published in peer-reviewed medical literature generally recognized by the relevant medical community, Physician Specialty Society recommendations and the views of Physicians practicing in relevant clinical areas and any other relevant factors.

‡ Indicated trademarks are the registered trademarks of their respective owners.

NOTICE: If the Patient’s health insurance contract contains language that differs from the BCBSLA Medical Policy definition noted above, the definition in the health insurance contract will be relied upon for specific coverage determinations.

NOTICE: Medical Policies are scientific based opinions, provided solely for coverage and informational purposes. Medical Policies should not be construed to suggest that the Company recommends, advocates, requires, encourages, or discourages any particular treatment, procedure, or service, or any particular course of treatment, procedure, or service.

NOTICE: Federal and State law, as well as contract language, including definitions and specific contract provisions/exclusions, take precedence over Medical Policy and must be considered first in determining eligibility for coverage.

